

19 Jun
13:00

Hypercyclic and mixing composition operators on $\mathcal{O}_M(\mathbb{R})$

Adam Przestacki

Adam Mickiewicz University

Dynamical properties of composition and weighted composition operators on various spaces of functions were intensively studied over the past decades by many authors e.g. by Bès, Bonet, Domaniński, Grosse-Erdmann, Jordá, Mortini, Shapiro.

During the talk I will focus on hypercyclicity and the mixing property of composition operators acting on the space of slowly increasing smooth functions $\mathcal{O}_M(\mathbb{R})$ well-known from the distribution theory. Recall that $\mathcal{O}_M(\mathbb{R})$ is given by

$$\mathcal{O}_M(\mathbb{R}) = \cap_{m=1}^{\infty} \cup_{n=1}^{\infty} \mathcal{O}_n^m(\mathbb{R}),$$

where

$$\mathcal{O}_n^m(\mathbb{R}) := \left\{ f \in C^m(\mathbb{R}) : |f|_{m,n} := \sup_{x \in \mathbb{R}, 0 \leq j \leq m} (1 + |x|^2)^{-n} |f^{(j)}(x)| < \infty \right\}.$$

I will present a characterization of all mixing composition operators C_ψ on $\mathcal{O}_M(\mathbb{R})$. Moreover I will show that this property is closely related to the solvability in $\mathcal{O}_M(\mathbb{R})$ of the Abel's functional equation, i.e. the problem to find $H \in \mathcal{O}_M(\mathbb{R})$ for a given symbol $\psi \in \mathcal{O}_M(\mathbb{R})$ which satisfies the equation

$$H(\psi(x)) = H(x) + 1.$$
